



ACCUEIL NOUVELLES ENTREPRISE SPORTS DIVERTISSEMENT INTERNATIONAL TECHNOLOGY

SCIENCES HEALTH CORONAVIRUS



Postes US News FR > Sciences

## Le graphène sur les surfaces de platine défie apparemment la loi de Coulomb

SCIENCE Zara Sciences about 14 hours ago



Étonnamment, le frottement entre la pointe d'un microscope à force atomique et les superstructures Moiré dépend de la vitesse à laquelle la pointe est déplacée sur la surface. Crédit : Département de physique, Université de Bâle et Scixel



Des chercheurs de Bâle et de Tel-Aviv ont découvert que la friction varie avec la vitesse dans des structures de graphène sur des surfaces de platine, défiant la loi de Coulomb qui stipule que la friction est indépendante de la vitesse dans le monde macro.

Materials made of single atomic layers are highly valued for their low-friction qualities, useful in reducing friction in hard disks or moving parts of satellites or space telescopes. Graphene, consisting of a single layer of carbon atoms arranged like a honeycomb, is a prime example and is under examination for its potential as a lubricating layer. Earlier studies showed that a graphene ribbon can glide almost friction-free across a gold surface.

### Surprising results with a rough surface

If graphene is applied to a platinum surface, it has a significant impact on the measurable friction forces. Now, physicists from the University of Basel and Tel Aviv University have reported in the journal *Nano Letters* that, in this instance, the friction depends on the speed at which the tip of an atomic force microscope (AFM; see box) is moved across the surface. This finding is surprising because friction does not depend on speed according to Coulomb's law, which applies in the macro world.

In conjunction with the platinum substrate, graphene no longer forms only the hexagonal honeycomb pattern of carbon atoms and instead forms superstructures known as Moiré superlattices. The surface is then no longer completely flat and exhibits a certain degree of roughness.

PROMOTED CONTENT

mgid ▶

"If we move the AFM tip across this slightly corrugated surface at low speed, we measure a weak and almost constant frictional force," explains Professor Ernst Meyer from the Swiss Nanoscience Institute and the Department of Physics at Basel University. "Above a certain threshold, however, the friction then increases with the speed of the AFM tip," adds first author Dr. Yiming Song. "The larger the Moiré superstructure, the lower the threshold at which the friction becomes speed-dependent."

The researchers found that there is greater resistance at the ridges of the Moiré superstructures during the movement of the tip. These ridges undergo elastic deformation due to the pushing tip before relaxing again when the pressure is sufficiently high. This effect results in greater frictional forces that increase with the speed of the tip. Simulations and an analytical model confirm the experimental findings obtained by this international team of researchers.

Reference: "Velocity Dependence of Moiré Friction" by Yiming Song, Xiang Gao, Antoine Hinaut, Sebastian Scherb, Shuyu Huang, Thilo Glatzel, Oded Hod, Michael Urbakh and Ernst Meyer, 30 November 2022, *Nano Letters*. DOI: 10.1021/acs.nanolett.2c03967

The article is in English

Tags: graphène sur les surfaces de platine défie apparemment la loi de Coulomb

PROMOTED CONTENT

mgid ▶

RELATED POSTS

This Video May Soon Be Banned. Watch Before It's Deleted

Altersfleck entfernen? Sie haben die Lösung zu Hause!

Beste Weltraumstrategie, um auf unbewohnte Planeten zu überleben

Makeup Artist Turns Ordinary Women Into Beauty Queens

8 Reasons Korean Women Don't Get Fat

Hottest Photos From Sports Events

30 Photos Of The Best Moments In Sports History

Is Your Partner Cheating? Look Out For These Common Signs

10 Mystic Forests Around The World

Historical Pics You Can't Unsee

Red Dragons Media

Zestradar

StyleFocus

Zestradar